



Thymic Epithelial Cell Medium (TyEpiCM) Catalog #3911

Product Description

Thymic Epithelial Cell Medium (TyEpiCM) is a complete medium designed for optimal growth of normal thymic epithelial cells *in vitro*. It is a sterile, liquid medium containing essential and non-essential amino acids, vitamins, organic and inorganic compounds, hormones, growth factors and trace minerals. The medium is serum-free. It is HEPES and bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5% CO₂/95% air. The medium is formulated (quantitatively and qualitatively) to provide a defined and optimally balanced nutritional environment that selectively promotes growth of normal thymic epithelial cells *in vitro*.

Components

TyEpiCM consists of 500 ml of basal medium, 5 ml of Thymic Epithelial Cell Growth Supplement (TyEpiCGS, Cat. #3962) and 5 ml of penicillin/streptomycin solution (P/S, Cat. #0503). **Note:** Due to bovine pituitary extract (BPE) in the growth supplement, the color may vary with different lots and formation of lipoproteins can result in precipitates. This does not affect the biological activity.

Product Use

TyEpiCM is for research use only. It is not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

Storage

Store the basal medium at 4°C, the TyEpiCGS and the P/S solution at -20°C. Protect from light.

Shipping

Dry ice.

Instructions for use

Thaw TyEpiCGS and P/S solution at 37°C. Gently tilt the TyEpiCGS tube several times to ensure the contents are completely dissolved before adding to the medium. Rinse the bottle and tubes with 70% ethanol and wipe to remove excess ethanol. In a sterile field, remove the cap, being careful not to touch the interior threads with fingers. Add TyEpiCGS and P/S solution to the medium and mix well. Since several components are light-labile, the medium should not be exposed to light for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.