

GoldNStart TaqGreen qPCR Master Mix (GTQMM)

Catalog #MB6018-1, 1 mL Catalog #MB6018-5, 5 mL Catalog #MB6018-50, 50 mL

Introduction

ScienCell's GoldNStart TaqGreen qPCR Master Mix (GTQMM) is a SYBR® Green dye-based qPCR master mix with a "hot-start" property. GTQMM is ideal for use in real-time quantitative PCR and DNA melt curve analysis. The 2X master mix contains SYBR® Green, dNTPs, Taq DNA polymerase, and an inert gold-color loading indicator (ScienCell, catalog #GQ300G) in a single tube. The "hot-start" property achieved through ScienCell's unique chemically modified Taq DNA polymerase provides maximal inhibition of primer dimer formation. The advanced buffer formulation provides superior specificity and efficiency with a wide linear dynamic range. The inert gold-color loading indicator allows for better visualization and tracking of sample loading in qPCR plates or tubes.

Kit Components

Catalog #MB6018-1

Cat #	Item	Quantity	Storage
MB6018a-1	GoldNStart TaqGreen qPCR Master Mix	1 mL	-20°C
MB6018b-1	Nuclease-free water	1 mL	4°C

Catalog #MB6018-5

Cat #	Item	Quantity	Storage
MB6018a-1	GoldNStart TaqGreen qPCR Master Mix	1 mL x 5	-20°C
MB6018b-1	Nuclease-free water	1 mL x 5	4°C

Catalog #MB6018-50

Cat #	Item	Quantity	Storage
MB6018a-10	GoldNStart TaqGreen qPCR Master Mix	10 mL x 5	-20°C
MB6018b-10	Nuclease-free water	10 mL x 5	4°C

Quality Control

The linear dynamic performance of GTQMM is verified with serially diluted DNA samples. DNase activity was NOT detected by incubating each component of GTQMM with single-stranded and double-stranded DNA at 37 °C for 24 hours.

Product Use

GTQMM is for research use only. It is not approved for human or animal use, or for application in clinical or in vitro diagnostic procedures.

Shipping and Storage

The product is shipped on dry ice. Upon receipt, store GoldNStart TaqGreen qPCR Master Mix (Cat #MB6018a) at -20°C in a manual defrost freezer and nuclease-free H₂O (Cat #MB6018b) at 4°C. Aliquot as needed. Avoid repeated freeze-and-thaw cycles.

Procedure

Important: Only use nuclease-free reagents in PCR amplification.

Note: This master mix does not contain a ROX passive reference dye. If the qPCR instrument being used has a "ROX passive reference dye" option, please deselect this option.

- 1. Thaw GoldNStart TagGreen qPCR Master Mix and place on ice.
- 2. Prepare 20 μL qPCR reactions in qPCR tubes or plates as shown in Table 1. For other reaction volume setup, scale up or down proportionally.

Table 1. Preparation of 20 µL qPCR reactions

Component	Volume	Final concentration
GoldNStart TaqGreen qPCR Master Mix	10 μL	1X
Template DNA	variable	-
Nuclease-free water	variable	-
Forward and reverse primers	variable	250 - 500 nM
Total volume per reaction	20 μL	-

- 3. Seal the qPCR reaction wells. Centrifuge the tubes or plates at 1,500X g for 15 seconds. For maximum reliability, replicates are recommended (minimum of 3).
- 4. Refer to Table 2 for a typical qPCR program setup. Adjust properly according to the optimized qPCR conditions for the reactions to run. Load the PCR tubes or plates into the qPCR instrument and run the program.

Table 2. A typical qPCR program setup

Step	Temperature	Time	Cycles
Taq DNA polymerase activation	95°C	10 min	1
Denaturation	95°C	20 sec	
Annealing	50 - 68°C	20 sec	30-45
Extension	72°C	20-45 sec	30-43
Data acquisition	Plate read		
Optional	Melting curv	e analysis	1
Hold	20°C	Indefinite	1

5. For data analysis, please refer to the data analysis software of the qPCR instrument being used.




Figure 1. Dynamic range and qPCR efficiency of GoldNStart TaqGreen qPCR Master Mix (ScienCell, Cat #MB6018). 10-fold serial dilution of human LDHA gene, ranging from 10^7 to 10^2 copies, were amplified in triplicates using GoldNStart TaqGreen qPCR Master Mix (ScienCell, Cat #MB6018), with no-template control (NTC) included.